

Bedrock Geologic Map of the Lowell (Iowa) 7.5' Quadrangle

LEGEND

CENOZOIC

QUATERNARY SYSTEM

Qu - **Undifferentiated Unconsolidated Sediments** - Consists of loose sand developed in loess, glacial till, and colluvium of varved blankets, and alluvial fill, silt, sand, and gravel. The total thickness of the Quaternary deposits varies between 0 and 10 to 20 ft. This unit is as much as 40 to 120 ft thick in the northern part of the mapping area. This unit is shown only on the cross-section, not on the map.

PALEOZOIC

CARBONIFEROUS SYSTEM

Pcl - **Sandstone and Shale** (lower Cherokee Group Lower Middle Pennsylvanian). Pennsylvanian units occur an erosional outlier reaching a thickness of up to 15 to 20 ft thick in the mapping area. The unit consists of thin-bedded, light to medium gray, fine to medium grained sandstone, and dark gray to black shale. The sandstone is typically sub- to medium grained, with coarse to medium beds, primary bedrock, and some thin-bedded. Some shales are calcareous to phosphatic, with minor coal and rare limestone. Lithologic patterns are commonly gray to dark brown dolomite, locally brecciated and sandy, with rare fossils. This mapping unit denotes the bedrock surface of the mapping area and is overlain by Quaternary sediments of Pennsylvanian origin. Outcrop of this mapping unit is found in the Skunk River Park area, several miles east of the Skunk River valley, and in the quarry.

MISSISSIPPIAN SUBSYSTEM

Mpsl - **Limestone, Sandstone, and Dolomite** (Pella or "St. Louis" Formations) Middle Upper Mississippian, Meramecian-Lower Chesterian. This unit is up to 120 to 140 ft thick in the mapping area. It is determined by limestone, sandstone, dolomite, chert, and locally abundant brachiopods, trilobites, and crinoids. The Pella and "St. Louis" Formations are typically sub- to medium grained, with coarse to medium beds, primary bedrock, and some thin-bedded. Some shales are calcareous to phosphatic, with minor coal and rare limestone. Lithologic patterns are commonly gray to dark brown dolomite, locally brecciated and sandy, with rare fossils. This mapping unit denotes the bedrock surface of the mapping area and is overlain by Quaternary sediments of Pennsylvanian origin. Outcrop of this mapping unit is found in the Skunk River Park area, several miles east of the Skunk River valley, and in the quarry.

Mws - **Limestone, Sandstone, and Dolomite** (Warsaw Formation) Upper Osagean. The Warsaw Formation varies in thickness reaching a maximum thickness of approximately 75 to 100 ft. This unit is generally divided into two major lithologic groups, a lower argillaceous dolomite sequence and an upper cherty dolomite sequence. The upper thick is typically light to medium gray, cherty, and locally dolomite with minor clay, sand, and some sparse fossils. The lower dolomite, sometimes referred to as the "light bed", is argillaceous to shaly, with scattered thin cherty partings. Warsaw limestone units occur locally in the Peoria, Iowa, with cherty beds, primary bedrock, and some thin-bedded. Some shales are calcareous to phosphatic, with minor coal and rare limestone. Lithologic patterns are commonly gray to dark brown dolomite, locally brecciated and sandy, with rare fossils. This mapping unit denotes the bedrock surface of the mapping area and is overlain by Quaternary sediments of Pennsylvanian origin. Outcrop of this mapping unit is found in the Skunk River Park area, several miles east of the Skunk River valley, and in the quarry.

Mk - **Limestone, Sandstone, and Dolomite** (Kinderhookian) Upper Osagean. The Kinderhookian sequence ranges in thickness from 20 to 60 ft in the mapping area. This unit is dominated by thin to medium grained, light to medium gray, cherty dolomite, with scattered thin cherty partings. The Kinderhookian sequence is generally divided into two major lithologic groups, a lower argillaceous dolomite sequence and an upper cherty dolomite sequence. The upper thick is typically light to medium gray, cherty, and locally dolomite with minor clay, sand, and some sparse fossils. The lower dolomite, sometimes referred to as the "light bed", is argillaceous to shaly, with scattered thin cherty partings. Kinderhookian limestone units occur locally in the Peoria, Iowa, with cherty beds, primary bedrock, and some thin-bedded. Some shales are calcareous to phosphatic, with minor coal and rare limestone. Lithologic patterns are commonly gray to dark brown dolomite, locally brecciated and sandy, with rare fossils. This mapping unit denotes the bedrock surface of the mapping area and is overlain by Quaternary sediments of Pennsylvanian origin. Outcrop of this mapping unit is found in the Skunk River Park area, several miles east of the Skunk River valley, and in the quarry.

Mb - **Dolomite, Limestone, and Chert** (Burlington Formation) Lower Osagean. The Burlington Formation can be up to 20 to 40 ft thick in the mapping area. This unit is subdivided into three members: the Burlington, Hight Creek, and Cedar Fork, characterized by distinct lithologic patterns. The Burlington Member is dominated by white to tan dolomite limestone displaying micritic and granitic textures, and includes a basal chert. The Hight Creek Member is characterized by dolomite with an interbedded sandstone and shale member referred to as the "waffle granitic", and thick beds of chert. A glauconitic zone marks the lower contact between the Burlington Member and Warsaw. The Cedar Fork Member is dominated by a thin bedded, micritic limestone. The Burlington Formation is a gray to white, crystalline limestone, locally calcareous and phosphatic, with scattered thin cherty partings. The Burlington Formation is a gray to white appearance. Occasional fish scales and graptolite are also observed in this member. Outcrop of the Burlington Formation is typically along the Skunk River and in the quarry and in rock quarries along the southeastern boundary of the mapping area. This unit is shown only on the cross-section, not on the map.



Mk - **Dolomite, Limestone, and Shale** (Kinderhookian) Lower Osagean. The Kinderhookian sequence ranges in thickness from 20 to 60 ft in the mapping area. This unit is dominated by thin to medium grained, light to medium gray, cherty dolomite, with scattered thin cherty partings. The Kinderhookian sequence is generally divided into two major lithologic groups, a lower argillaceous dolomite sequence and an upper cherty dolomite sequence. The upper thick is typically light to medium gray, cherty, and locally dolomite with minor clay, sand, and some sparse fossils. The lower dolomite, sometimes referred to as the "light bed", is argillaceous to shaly, with scattered thin cherty partings. Kinderhookian limestone units occur locally in the Peoria, Iowa, with cherty beds, primary bedrock, and some thin-bedded. Some shales are calcareous to phosphatic, with minor coal and rare limestone. Lithologic patterns are commonly gray to dark brown dolomite, locally brecciated and sandy, with rare fossils. This mapping unit denotes the bedrock surface of the mapping area and is overlain by Quaternary sediments of Pennsylvanian origin. Outcrop of this mapping unit is found in the Skunk River Park area, several miles east of the Skunk River valley, and in the quarry.

Der - **Siltstone and Shale** (English River Formation) Upper Devonian, lower to upper Famennian. The English River Formation is up to 6 to 10 ft thick in the mapping area. This unit is dominated by gray to olive green siltstone with sparse to abundant fusulinid fossils. Lithology and thickness varies, especially in the upper beds, with siltstone to shaly sandstone. This unit appears only on the cross-section and not on the map.

Dss - **Shale** (Saverton Shale Formation) Upper Devonian, lower to upper Famennian. The Saverton Shale Formation can be up to 40 to 100 ft thick in the mapping area. This unit is dominated by green gray shale, commonly brecciated with sparse to abundant fusulinid fossils. This unit appears only in the cross-section and not on the map.

OTHER FEATURES

- New drill holes for this map project
- ✕ Bedrock outcrop
- ✕ ICG GEORAN data points - records available at www.geogran.iastate.edu
- Quarries
- Road
- W24902 Well used for geologic cross-section
- Bedrock Hillshade - shades of gray show the bedrock surface as it would be illuminated by an artificial light source from the SW direction



BEDROCK GEOLOGIC MAP OF THE LOWELL 7.5' QUADRANGLE, DES MOINES, HENRY, AND LEE COUNTIES, IOWA

Open File Map W-14-17-5
June 2017

Ryan Clark, Huihao Liu, Stephanie Tassier-Surine and Phil Kerr
Iowa Geological Survey, IIHR-Hydroscience & Engineering, University of Iowa, Iowa City, Iowa

Iowa Geological Survey, Robert D. Liles, State Geologist

Supported in part by the U.S. Geological Survey
Cooperative Agreement Number G14AC00103
National Cooperative Geologic Mapping Program (NCEM/NCGMP)
Contracted under award by the Department of Energy

ACKNOWLEDGMENTS

Special thanks to Peter Smith of Coalbrook Construction Co., Oklahoma Materials, Chuck Mason of Dumb Stone, Ryan Gearty of the Burlington Data Department, and Jeffrey of the Iowa Army Ammunition Plant Keady Collection. Above: Ronalda Doss Wajpalee, Dusan Golea, Zhaa McAuley, Jan Kambouch, and Doug Forman are thanked for allowing us access to their property. Zhaa Forman, Director of the Iowa Geological Survey (IGS) and University of Iowa Geology Department, is thanked for his support and assistance in the field. The following individuals are thanked for their assistance: James Harbeck and Dan Penland, US Department of Earth and Environmental Sciences students by producing descriptive logs of water well drilling records; Cliff Metchalk, Lyle W. Kott, and Robert M. Kott, Iowa Geological Survey; and the following: Robert M. Kott, Robert M. Kott, and Robert M. Kott. Outcrop of the Burlington Formation was observed at the Skunk River Park area, several miles east of the Skunk River valley, and in the quarry. Outcrop of the Burlington Formation was observed at the Skunk River Park area, several miles east of the Skunk River valley, and in the quarry. Outcrop of the Burlington Formation was observed at the Skunk River Park area, several miles east of the Skunk River valley, and in the quarry. Outcrop of the Burlington Formation was observed at the Skunk River Park area, several miles east of the Skunk River valley, and in the quarry.

Introduction to the Bedrock Geologic Map of the Lowell 7.5' Quadrangle, Des Moines, Henry, and Lee Counties, Iowa

The Lowell Quadrangle lies within the Southern Iowa Drift Plain landform region (Prior, 1991). This area hosts glacial deposits over 500,000 years old containing a thick till package mantled by loess draped over upland hill slopes. Numerous rills, creeks, and rivers branch out across the landscape shaping the local glacial deposits into steeply rolling hills and valleys.

The thickness of Quaternary materials overlying the bedrock surface varies widely across the quadrangle ranging from 0 to 18 m (0 to 60 ft), reaching a maximum thickness of 64 m (210 ft) in the northern part of the mapping area. Shallow bedrock information from the soil surveys in Des Moines, Henry, and Lee counties (Brown, 1983; Scholten, 1985; and Lockridge, 1979) was used for identifying potential bedrock outcrop locations during field mapping activities. Bedrock outcrops were found mostly along the Skunk River and its tributary creeks like Mud Creek and Cedar Creek, primarily in the eastern half of the quadrangle. Outcrops exposed are comprised of Pella and "St. Louis", Warsaw, and Keokuk formation rocks with one outcrop of Pennsylvanian bedrock. Subsurface information was mostly derived from the analysis of water well cutting samples stored at the Iowa Geological Survey (IGS). Lithologic and stratigraphic information from these samples are reported in the online GeoSam database of the IGS. Geologic information from 17 outcrops, and more than 200 private and public wells within the Lowell Quadrangle and surrounding area were used for bedrock geologic mapping purposes.

The Mississippian System (now Subsystem) was historically proposed for the succession of strata exposed in the Mississippi River Valley between Burlington, Iowa and southern Illinois. Therefore, the bedrock exposures in southeastern Iowa take on a special significance as they comprise part of the historic "body stratotype" on which the concept of the Mississippian System was defined and based (Witzke et al., 2002). The Mississippian had been primarily a North American chronostratigraphic label roughly synonymous with the Lower Carboniferous of the Old World. After approval by the Subcommittee on Carboniferous Stratigraphy in 1999 and ratification by the International Union of Geological Sciences and the International Commission on Stratigraphy in 2000, the Carboniferous System was officially subdivided into lower and upper subsystems, the Mississippian and Pennsylvanian, respectively. As such, the Mississippian now has meaning and application as a major subdivision of geologic time not only in North America, but as a globally defined subsystem of geologic time in the Lowell Quadrangle and the surrounding area provide a significant historic reference for the Mississippian as a whole.


The contention that is the Mississippian in Iowa has been the subject of curiosity for many previous workers. Owen (1852) and Hall (1857) were the first to recognize that the abundant bedrock exposures in southeastern Iowa likely correlated with those observed farther down the Mississippi River Valley. Decades later, Van Tuyl (1923) took on the ambitious task of correlating all of the Mississippian units across Iowa. Many of his lithologic interpretations were questionable, however, the correlations were and continue to be, subject to revision as later workers attempted to piece the Mississippian into the global stratigraphic framework. Harris and Parker (1964) provided inspirational insights into the structural context of southeastern Iowa by identifying a series of north-south trending anticlines that were later found to be superimposed on the larger northeast-southwest trending structural feature known as the Mississippi Arch (Witzke et al., 1990, p. 5). Many questions remain regarding the stratigraphic correlations within the Mississippian such as whether the "St. Louis" Formation in Iowa truly belongs in the St. Louis Formation or should be of the upper member be reassigned to the St. Genevieve Formation; whether the Prospect Hill Formation is an offshoot of the Hannibal Formation in Missouri and Illinois; and whether the Warsaw Formation is correlative to the McCraney in Illinois or if it should become a new stratigraphic interval (as proposed by Witzke et al., 2002). Although the Mississippian bedrock in southeastern Iowa is no longer a widely used aquifer due to low yields and locally poor water quality, many of the bedrock units are highly desirable sources of aggregate, thus necessitating the continued effort to gain a better understanding of the local and regional stratigraphic characteristics and relationships of the Mississippian System in southeastern Iowa.

The mapping area is dominated by bedrock of the Mississippian System that was deposited in a variety of marine environments from the late Kinderhookian to early Chesterian, approximately 355 to 330 million years ago (Ogg et al., 2008). Kinderhookian strata represent a sequence of interbedded carbonates and siltstones that unconformably underlie the Burlington Formation (early Osagean) and are not exposed at the bedrock surface within the mapping area. The Burlington, Keokuk, and Warsaw formations (collectively the Augustan Group of Witzke et al., 2010) represent a relatively conformable package of marine rocks deposited during the Osagean transgressive-regressive (T-R) cycle. Interpreted as part of the central middle shelf of the Osagean sea that transgressed toward the northwest and the Transcontinental Arch, the Burlington Formation rocks were deposited across a vast subtidal epicontinental shelf that stretched from Illinois and Iowa into central Kansas and Oklahoma (Lane, 1978; Witzke et al., 1990, p. 55). The Keokuk and Warsaw formations represent the regressive phase of the Osagean T-R cycle punctuated by a stark unconformity below the overlapping Pella and "St. Louis" formations, regionally displaying up to 40 m (130 ft) of erosional relief (Witzke et al., 2002). The Pella and "St. Louis" formations are mapped as one unit due to their stratigraphic complexity and questionable correlation to the type sections in Missouri and Illinois (Witzke et al., 1990, p. 23). The Pella and "St. Louis" formations were deposited in a near-shore environment as evidenced by mudflat facies rocks, evaporites and associated collapse breccias, and increased terrigenous sandstone deposits, with periods of brackish and/or lacustrine deposition interpreted from coal deposits and root casts (Witzke et al., 2002). Pennsylvanian bedrock units of the lower Cherokee Group occur an erosional outlier and are not well exposed in the mapping area. For a more detailed description of the lithologic units and further discussion of mapping methodologies, please refer to the accompanying Summary Report.


Base map from Iowa DOT road map layers 2006. Bedrock topography raster created internally for this map project.
New Geological Survey digital cartographic file Lowell_BedrockGeology.mxd, version 05/27/14 09:52:23
Map projection and coordinate system based on Universal Transverse Mercator (UTM) Zone 10N, datum NAD83.
The map and cross-section are based on interpretations of the best available information at the time of mapping. Map interpretations are not a substitute for detailed site specific studies.
Research supported by the U.S. Geological Survey, National Cooperative Geologic Mapping Program, under USGS award number G14AC00103. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

Adjacent 7.5' Quadrangles

MT PLEASANT IOWA	NEW LONDON IOWA	PLEASANT GROVE IOWA
SALEM IOWA	LOWELL IOWA	DAVILE IOWA
DONNELSON IOWA	WEST POINT IOWA	FORT WASHINGTON IOWA



GEOLOGIC CROSS-SECTION A-A'



Vertical exaggeration=10x