

**SUMMARY REPORT OF THE  
SURFICIAL GEOLOGIC MAP OF  
THE DANVILLE 7.5' QUADRANGLE, DES MOINES,  
HENRY, AND LEE COUNTIES, IOWA**

**Iowa Geological Survey  
Open File Map OFM-17-8  
June 2017**

Stephanie Tassier-Surine, Phil Kerr, Ryan Clark, and Huaibao Liu

Iowa Geological Survey, IIHR-Hydrosience & Engineering, University of Iowa, Iowa City, Iowa



Iowa Geological Survey, Robert D. Libra, State Geologist

Supported in part by the U.S. Geological Survey  
Cooperative Agreement Number G16AC00193  
National Cooperative Geologic Mapping Program (STATEMAP)  
Completed under contract with the Iowa Department of Natural Resources



## INTRODUCTION

The ‘Surficial Geologic Map of the Danville 7.5’ Quadrangle, Des Moines, Henry, and Lee Counties, Iowa’ is part of the first phase of surficial mapping in southeast Iowa. A bedrock geologic map was also produced in conjunction with the surficial geologic map (Clark et al., 2017). The Danville Quadrangle covers an area from 40° 45’ to 40° 52’ 30” N latitude and 91° 22’ 30” to 91° 15’ W longitude. The Danville Quadrangle is located on the Southern Iowa Drift Plain landform region (Prior and Kohrt, 2006). The map area is dominated by loess mantled till plains in the uplands, and glacial outwash and finer-grained alluvial deposits within the Skunk River valley and its tributaries. Stratigraphically, this area contains Illinoian age glacial deposits, which are only present in a small area of southeastern Iowa. The terminal moraine for the Illinoian glacial advance is three to four miles west of the mapping area. The thickness of Quaternary materials varies widely across the quadrangle generally ranging from 0 to 18 m (0-60 ft), and reaching a maximum thickness of 64 m (210 ft) in the northern part of the mapping area. Bedrock outcrops are located primarily in the southern half of the quadrangle, along the Skunk River and its tributaries including Cedar, Long, and Deeds creeks.

Mapping the Lowell and Danville quadrangles provides the first study of the regional Quaternary stratigraphy of southeast Iowa in almost 40 years since Hallberg (1980a,b) established the stratigraphy for the Illinoian and Pre-Illinoian glacial advances in eastern and southeastern Iowa. The majority of the drill cores and outcrops for those studies were north and east of the Danville Quadrangle. Additional data available since that time (LiDAR, DEMs, and digital soil surveys), have allowed for the refinement of the Illinoian boundary and greater detail in mapping the valleys. The only other surficial map of the area consists of the Des Moines 4° x 6° Quadrangle at a scale of 1:1,000,000 (Hallberg et al., 1991). Several Iowa Geological Survey (IGS) field trip guidebooks outline the Pleistocene, Devonian, and Mississippian stratigraphy (Witzke et al., 2002; Witzke and Tassier-Surine, 2001), but their focus was on the area near Burlington (to the east).

## PURPOSE

Detailed geologic mapping in southeast Iowa was completed as part of the IGS’s ongoing participation in the United States Geological Survey (USGS) STATEMAP Program. Mapping was completed as part of the IGS Developing Areas and Impaired Watershed mapping initiatives and provides comprehensive surficial and bedrock geologic information. These maps are the basis for further development of derivative datasets and map products for use by local, county and state decision-makers. An increased demand for groundwater resources in the region, new research into the Lower Skunk River watershed, development of additional aggregate resources, and expanding urban areas led to the selection of southeast Iowa as the next target for geologic mapping by the Iowa State Mapping Advisory Committee (SMAC). Key societal concerns that can be aided by this mapping project include watershed management, groundwater quantity and quality assessment, flood mitigation, aggregate resource protection, and land use planning and development.

Bedrock mapping efforts were successful in subdividing the Mississippian Augusta Group mapping unit used by Witzke and others (2010) into formations and in better identifying Pennsylvanian outliers. From a Quaternary perspective, characterizing and identifying the extent of the Illinoian glacial advance and establishing the loess thickness were primary objectives of the surficial map. The map area also includes the Skunk River, which carried outwash from the Wisconsin age Des Moines Lobe glacial

advance. Mapping has also helped to further delineate alluvial terraces, as well as identify and characterize sand and gravel resources associated with glacial outwash. Combining the bedrock and surficial map information is allowing stakeholders to address key questions related to shallow rock areas, groundwater protection, water supply concerns, and aggregate resource potential and protection.

## **QUATERNARY HISTORY AND REGIONAL SETTING**

The glacial history of Iowa began more than two million years ago, as at least seven episodes of Pre-Illinoian glaciation occurred between approximately 2.6 and 0.5 million years ago (Boellstorff, 1978a,b; Hallberg, 1980a). Early researchers believed there were only two episodes of Pre-Illinoian glaciation in Iowa. Later regional studies determined that at least seven episodes of Pre-Illinoian glaciation had occurred and led to the abandonment of the classic glacial and interglacial terminology: Kansan, Aftonian and Nebraskan (Boellstorff, 1978a,b; Hallberg, 1980a, 1986). Hallberg (1980a,b, 1986) undertook a regional scale project in east-central Iowa that involved detailed outcrop and subsurface investigations, including extensive laboratory work and synthesis of previous studies. Hallberg's study marked a shift from the use of time-stratigraphic terms and resulted in the development of a lithostratigraphic framework for Pre-Illinoian till. In east-central Iowa, Hallberg formally classified the units into two formations on the basis of differences in clay mineralogy: the Alburnett Formation (several undifferentiated members) and the younger Wolf Creek Formation (including the Winthrop, Aurora and Hickory Hills members). Both formations are composed predominantly of till deposits, but other materials are present. Paleosols are formed in the upper part of these till units.

A limited area of southeastern Iowa was glaciated during the Illinois Episode, around 300,000-130,000 years ago (Hallberg, 1980b). The Danville Quadrangle was glaciated during this time. The Illinoian till was deposited by the advancing Lake Michigan Lobe which moved across western Illinois into Iowa from the northeast (Leverett, 1899; Wickham, 1980). The Lake Michigan Lobe incorporated Paleozoic bedrock materials from the Lake Michigan Basin which are distinguished by both the clay mineralogy of the matrix as well as the pebbles and clasts (Lineback, 1980; Wickham, 1980). Coal fragments were commonly identified in core; this is unique to Illinoian age glacial till in Iowa. Hallberg (1980b) defined the formal stratigraphic nomenclature in Iowa. The only Illinoian Episode till present in Iowa is the Glasford Formation Kellerville Till Member (Willman and Frye, 1970). Following the Illinoian glaciation, this area underwent landscape development and erosion until the Wisconsin Episode loess began to be deposited. Illinoian till is only exposed in drainages and relatively steep sideslopes.

In eastern Iowa, the highly eroded and dissected Illinoian and Pre-Illinoian upland and older terraces are mantled by two Wisconsin loesses. The older Pisgah Formation is thin and includes loess and related slope sediments that have been altered by colluvial hillslope processes. The unit is characterized by the presence of a weakly developed soil recognized as the Farmdale Geosol. It is not uncommon to see the Farmdale developed throughout the Pisgah Formation and into the underlying older Sangamon Paleosol. The Pisgah loess was most likely deposited on the eastern Iowa landscape from 30,000 to 24,000 years ago (Bettis, 1989) and is typically buried by Peoria Formation loess. The Peoria Formation loess accumulated on stable landsurfaces in eastern Iowa from 25,000 to 21,000 years ago. Peoria Formation eolian materials mantle the upland till units and are present on the Wisconsin outwash terraces. On the uplands, the Peoria Formation is a uniform silt loam; in the valleys the silt commonly grades downward to fine sand. The loess deposits in the mapping area are relatively thin, generally less than 4 meters (12 ft).

The Skunk River deposited coarse sand and gravel associated with glacial outwash (Noah Creek Formation) of the Des Moines Lobe during the Wisconsin Episode. The sand and gravel is much thinner (typically less than 6 m) than in many other rivers in Iowa that carried Wisconsin outwash. These deposits are typically overlain by Holocene alluvium. Hudson age deposits are associated with fine-grained alluvial, organic, and colluvial sediments and include the DeForest Formation which is subdivided into the Camp Creek, Roberts Creek, and Gunder members. These deposits are present in valleys and upland drainages throughout the map area. The Holocene low terrace deposits occupy the active channel belt of the Skunk River. Both an intermediate and high Holocene terrace are present along the Skunk River and may be several meters above the modern floodplain. Due to the difficulty of differentiating these terraces where only one was present, they were combined into one mapping unit.

## **METHODS**

Numerous existing sources of geologic information were utilized in the production of the surficial and bedrock geologic maps of the Danville Quadrangle including subsurface information, USDA NRCS soil survey data, aerial photography, DEM's, satellite imagery, landform characteristics, and LiDAR. Where available, engineering borings from public utilities, the Iowa Department of Transportation, and monitoring well records of the USGS were used. Subsurface lithologic and stratigraphic information was mostly derived from analysis of water well cutting samples repositied at the IGS and stored in the IGS online GeoSam database. Over 260 public and private wells in GeoSam, including strip logs, were reviewed for lithology, stratigraphy and locational accuracy, and updated where needed. NRCS digitized soils data (Brown, 1983; Seaholm, 1985; and Lockridge, 1979) provided information regarding shallow rock areas, helped to guide valley mapping units, and defined slope areas where glacial till is exposed. Bedrock mappers also used the digital soil surveys to help delineate areas of shallow rock outcrop prior to field reconnaissance. New geologic information was obtained from field investigations of 31 outcrops (including quarry exposures) and logging of well cutting samples for 66 unstudied wells totaling 17,416 feet. Quaternary geologists worked with a contract driller and utilized the IGS Giddings probe to drill a mix of solid stem and continuous core holes. Five new drill holes totaling 101 feet were completed in the quadrangle to characterize the Quaternary sediments and establish unit thickness. Samples have been submitted to the Quaternary Materials Lab at the University of Iowa's Earth and Environmental Sciences Department for grain-size analysis. All results are expected by August, 2017. Project geologists combined information from the sources listed above to delineate surficial geologic mapping units at 1:24,000 scale for the Danville Quadrangle. IGS mappers used ArcGIS and on-screen digitizing techniques developed during previous STATEMAP projects. The final map entitled 'Surficial Geologic Map of the Danville 7.5' Quadrangle, Des Moines, Henry, and Lee Counties, Iowa' will be available as a shapefile in the Iowa Department of Natural Resources NRGIS library, as a PDF file on the IGS Publications website, and will be submitted to the USGS National Geologic Map Database. This Summary Report is also available as a PDF file on the IGS Publications website.

## **STRATIGRAPHIC FRAMEWORK FOR SOUTHEAST IOWA**

The stratigraphic framework for southeast Iowa was established by Hallberg (1980a,b) nearly 40 years ago. Surficial deposits in the map area are composed of seven formations (youngest to oldest): Hudson DeForest; Wisconsin Peoria, Pisgah, and Noah Creek; Illioian Glasford; and Pre-Illinoian Wolf Creek and Alburnett. Hudson age deposits associated with fine-grained alluvial, organic, and colluvial

sediments include the DeForest Formation which is subdivided into the Camp Creek, Roberts Creek, Gunder, and Corrington members. The Noah Creek Formation includes coarse sand and gravel associated with outwash from the Des Moines Lobe. Loess deposits include both Peoria and Pisgah formation silt that are present mantling the upland till units and are found intermittently on Holocene terraces. The only Illinoian till unit present in Iowa is the Glasford Formation Kellerville Member. The primary work on these deposits was completed to the north of the current mapping area. Although numerous drill holes in the previous study suggested a supraglacial facies may be present, limited evidence was found in the current study. Illinoian glacial sediments are found throughout the mapping area and consists of uniform dense till. Pre-Illinoian glacial deposits are not exposed at the surface, but are present at depth and consist of two formations: the younger Wolf Creek Formation and the Alburnett Formation. The Wolf Creek Formation is divided into the Winthrop, Aurora, and Hickory Hills members (oldest to youngest). The Alburnett Formation consists of several “undifferentiated” members.

Six bedrock mapping units (Pennsylvanian Lower Cherokee Group; and the Mississippian Pella and “St. Louis”, Warsaw, Keokuk, Burlington, and Kinderhookian formations) are exposed at the bedrock surface in the Danville Quadrangle. The Mississippian Pella and “St Louis” Formation and the Pennsylvanian Lower Cherokee Group comprise the bedrock in most of the map area, especially in the upland areas. The other Mississippian units occur within the bedrock valleys and tributaries. Twenty-six rock outcrops and five quarries are located in the map area and were investigated in the field. Bedrock exposures or rock present within one to two meters (7 ft) of the land surface are designated as ‘Qbr’ on the map. Specific bedrock units are shown on the cross-section and defined in the legend. For detailed bedrock information see The Bedrock Geologic Map of the Danville 7.5’ Quadrangle (Clark et al., 2017).

Recent studies and mapping indicate that the map area encompasses a complex suite of depositional landforms and sediment sequences related to glaciations, alluviation, subaerial erosion, and wind-blown transport. To map diverse landscapes at 1:100,000 scale, we have selected the most comprehensive mapping strategy- a landform sediment assemblage (LSA) approach. Various landforms are the result of specific processes at work in the geologic system. Landforms typically have similar relief, stratigraphic and sedimentologic characteristics. Recognition of the genetic relationship among landforms and their underlying sediment sequences allows one to generalize and map complex glacial terrains over areas of large extent (Sugden and John, 1976; Eyles and Menzies, 1983). Bettis and others (1999) found that LSA mapping concepts were extremely useful in overcoming the difficulties of mapping in large valleys and noted that LSA’s provided a unique opportunity to associate landforms with their underlying sediment packages. Eight landform sediment assemblage units were identified in the map area utilizing aerial imagery, topographic expression, digitized soils, LiDar and existing and new subsurface geologic boring information. The following is a description of each landform sediment assemblage listed in order of episode:

### **HUDSON EPISODE**

**Qal - Alluvium** (DeForest Formation-Undifferentiated) Variable thickness of less than 1 to 5 m (3-16 ft) of very dark gray to brown, noncalcareous to calcareous, stratified silty clay loam, clay loam, loam to sandy loam alluvium and colluvium in stream valleys, on hill slopes and in closed depressions. May overlie Glasford Formation glacial till, Peoria Formation loess or eolian sand, or Noah Creek Formation sand and gravel. Associated with low-relief modern floodplain, closed depressions, modern drainageways or toeslope positions on the landscape. Seasonal high water table and potential for frequent flooding.

**Qallt - Low Terrace** (DeForest Formation-Camp Creek and Roberts Creek members) Variable thickness of less than 1 to 5 m (3-16 ft) of very dark gray to brown, noncalcareous, stratified silty clay loam, loam, or clay loam, associated with the modern channel belt of the Skunk River valley. Overlies Noah Creek Formation sand and gravel. Occupies the lowest position on the floodplain ie. modern channel belts. Seasonal high water table and frequent flooding potential.

**Qali-ht - Intermediate-High Terrace** (DeForest Formation-Gunder Member) Variable thickness of less than 1 to 5 m (3-16 ft) of very dark gray to brown, noncalcareous, silty clay loam to loam alluvium or colluvium. Overlies Noah Creek Formation sand and gravel along the Skunk River. Occupies terrace and valley margin positions 1 to 2 m (3-7 ft) above the modern floodplain. Two terrace levels are present in some areas. Seasonal high water table and low to moderate flooding potential.

### **WISCONSIN EPISODE**

**Qnw - Sand and Gravel** (Noah Creek Formation) Generally 3 to 9 m (10-30 ft), but may be up to 15 m (50 ft), of yellowish brown to gray, poorly to well sorted, massive to well stratified, coarse to fine feldspathic quartz sand, pebbly sand and gravel with few intervening layers of silty clay. This unit is buried by younger Hudson-age alluvial deposits associated with the Skunk River valley and encompasses deposits that accumulated in river valleys during the Wisconsin Episode. This unit is shown only on the cross-section.

**Qps-gla - Loess** (Peoria Formation-silt facies) Generally 2 to 5 m (7-15 ft) of yellowish to grayish brown, massive, jointed, calcareous or noncalcareous silt loam to silty clay loam. May overlie a grayish brown to olive gray silty clay loam to silty clay (Pisgah Formation- eroded Farmdale Geosol) which is less than 1.5 m (5 ft) thick. The Pisgah Formation is in the same stratigraphic position as the Roxanna Silt which is mapped in Illinois. The Farmdale Geosol may be welded to an older Sangamon Geosol developed in loamy glacial till of the Glasford Formation. This mapping unit encompasses upland divides, ridgetops and convex sideslopes. Well to somewhat poorly drained landscape.

### **ILLINOIS EPISODE**

**Qgla - Till** (Glasford Formation) Generally 3 to 10 m (10-33 ft) of very dense, massive, fractured, loamy glacial till of the Illinoian Glasford Formation with or without a thin loess mantle (Peoria Formation- less than 2 m) and intervening clayey Farmdale/Sangamon Geosol. Overlies the Yarmouth Paleosol formed in Pre-Illinoian till. This mapping unit encompasses narrowly dissected interfluvial and side slopes, and side valley slopes. Drainage is variable from well drained to poorly drained.

### **PRE-ILLINOIS EPISODE**

**Qwa3 - Till** (Wolf Creek or Alburnett formations) Generally 10 to 35 m (33-115 ft) of very dense, massive, fractured, loamy glacial till of the Wolf Creek or Alburnett formations. The Yarmouth Paleosol is formed in this unit, and it is overlain by Illinoian till. Drainage is variable from well drained to poorly drained. This unit is shown only on the cross-section.

**Qbr - Loamy Sediments Shallow to Dolomite, Limestone, Shale and Sandstone** (DeForest, Noah Creek, Peoria, and Glasford formations) - 1 to 2 m (3-7 ft) of yellowish brown to gray, massive to weakly stratified, well to poorly sorted loamy, sandy and silty sediments that overlie the Pennsylvanian or Mississippian bedrock surface. All areas of bedrock outcrop or shallow to bedrock soils are shown in red on the map, regardless of the bedrock mapping unit. Bedrock units are shown on the cross-section and may be identified on the bedrock map of the Danville Quadrangle.

## ACKNOWLEDGEMENTS

Special thanks to the landowners who allowed access to their properties for drilling: Herb Grinstead, Dennis Harwood, Chris Lee (Director) and Jeff Breuer of the Des Moines County Conservation Board, Dick Moore, Kory Moore, John Pearson and Jim Jansen of the Iowa Department of Natural Resources (IDNR), Don Schnupper, and Ron Stevenson. Drilling was provided under contract with Cahoy Well and Pump Service; special thanks to driller Mark Claassen. Thanks also to Don Smith of Cessford Construction Co. (Oldcastle Materials), Chuck Manson of Douds Stone, Ryan Gourley of the Burlington Parks Department, and Joe Haffner of the Iowa Army Ammunition Plant. Zachary Demanett of the Iowa Geological Survey (IGS) and University of Iowa (UI) students Samantha Moser, Ryan McKeon, and Gia DeBartolo prepared well cutting samples for stratigraphic logging. New subsurface geologic data was generated by Tanner Hartsock and Diar Ibrahim, UI Department of Earth and Environmental Sciences students, by producing descriptive logs of water well drilling samples. UI students Nick Lefler and Katie Grob helped with checking well locations and data management. Zachary Demanett and Tanner Hartsock also assisted with field work. Thanks also to Rick Langel (IGS) for managing the Iowa geologic sampling database (GeoSam). Also, special thanks to Deborah Quade (IDNR) and Art Bettis, UI Department of Earth and Environmental Sciences, for numerous valuable discussions regarding the geology of southeast Iowa. Casey Kohrt and Chris Kahle of IDNR and Andrew Roers of IHR Hydroscience & Engineering provided GIS technical help. Administrative support was provided by Megan Delaney, Rosemary Tiwari, Teresa Gaffey, Angi Roemerman, and Carmen Langel.

## REFERENCES

- Bettis, E.A., III, 1989, Late Quaternary history of the Iowa River Valley in the Coralville Lake area in Plocher, O.W., Geologic Reconnaissance of the Coralville Lake area: Geological Society of Iowa Guidebook 51, p. 93-100.
- Bettis, E.A. III, Hajic, E.R., and Quade, D.J., 1999, Geologic mapping of large valleys in glaciated regions: The use of landform and landscape sediment assemblages for multi-use maps: Geological Society of America Abstracts with Programs, 33<sup>rd</sup> Annual Meeting North-Central Section April 1999, Champaign, Illinois, v. 31, no. 5, abstract no. 04164.
- Boellstorff, J., 1978a, North American Pleistocene Stages reconsidered in light of probable Pliocene-Pleistocene continental glaciation: *Science*, v. 202, p. 305-307.
- Boellstorff, J., 1978b, Chronology of some late Cenozoic deposits from the central United States and the ice ages: *Transactions of the Nebraska Academy of Science*, v. 6, p. 35-49.
- Brown, M.D., 1983, Soil Survey of Des Moines County, Iowa: U.S. Department of Agriculture, Soil Conservation Service, 199 p., 66 map sheets.
- Clark, R., Liu, H., Tassier-Surine, S., and Kerr, P., 2017, Bedrock Geologic Map of the Danville 7.5' Quadrangle, Des Moines, Henry, and Lee Counties, Iowa: Iowa Geological Survey, Open File Map OFM-17-7, 1:24,000 scale map sheet.
- Curry, B.B. and Grimley, D.A., 2006, Provenance, age and environment of mid-Wisconsinan slackwater lake sediment in the St. Louis Metro East area, USA: *Quaternary Research*, v. 65, p. 108-122.
- Esling, S.P., 1984, Quaternary stratigraphy of the lower Iowa and Cedar River valleys, southeast Iowa: University of Iowa, Iowa City, unpublished PhD Dissertation, 451 p.
- Eyles, N. and Menzies, J., 1983, The subglacial landsystem *in* Eyles, N., ed., *Glacial geology-An introduction for engineers and earth scientists*: Oxford, Pergamon, p. 19-70.
- Hallberg, G.R., 1980a, Pleistocene stratigraphy in east-central Iowa: Iowa Geological Survey Technical Information Series, v. 10, 168 p.
- Hallberg, G.R., ed., 1980b, Illinoian and Pre-Illinoian stratigraphy of southeast Iowa and adjacent Illinois: Iowa Geological Survey Technical Information Series, v. 11, 206 p.
- Hallberg, G.R., 1986, Pre-Wisconsin glacial stratigraphy of the central plains region in Iowa, Nebraska, Kansas, and Missouri *in* Sibrava, V., Bowen, D.Q., and Richmond, G.M., eds., *Quaternary Glaciations in the Northern Hemisphere: Quaternary Science Reviews*, v. 5, p. 11-15.
- Hallberg, G.R., Lineback, J.A., Mickelson, D.M., Knox, J.C., Goebel, J.E., Hobbs, H.C., Whitfield, J.W., Ward, R.A., Boellstorff, J.D., and Swinehart, J.B., 1991, Quaternary geologic map of the Des Moines 4° x 6° quadrangle, United States: U.S. Geological Survey, Miscellaneous Investigations Series, Map I-1420, 1:100,000 scale map sheet.
- Leverett, F., 1899, The Illinois glacial lobe: U.S. Geological Survey, Monograph 38, 817 p.
- Lineback, J.A., 1980, The Glasford Formation of western Illinois in Hallberg, G.R., Illinoian and Pre-Illinoian stratigraphy of southeast Iowa and adjacent Illinois: Iowa Geological Survey Technical Information Series, v. 11, p. 181-184.
- Lockridge, L.D., 1979, Soil Survey of Lee County, Iowa: U.S. Department of Agriculture, Soil Conservation Service, 188 p., 85 map sheets.
- Prior, J.C. and Kohrt, C.J., 2006, The Landform Regions of Iowa: Iowa Geological Survey, digital map, available on IDNR NRGIS Library.
- Seaholm, J.E., 1985, Soil Survey of Henry County, Iowa: U.S. Department of Agriculture, Soil Conservation Service, 259 p., 63 map sheets.
- Sugden, D.E., and John, B.S., 1976, *Glaciers and Landscape*: New York, John Wiley & Sons, 376 p.
- Wickham, J.T., 1980, Status of the Kellerville Till Member in western Illinois in Hallberg, G.R., ed., 1980: Illinoian and Pre-Illinoian stratigraphy of southeast Iowa and adjacent Illinois, Iowa Geological Survey Technical Information Series, v. 11, p. 151-180.



- Willman, H.B. and Frye, J.C., 1970, Pleistocene stratigraphy of Illinois: Illinois State Geological Survey Bulletin 94, 204 p.
- Witzke, B.J. and Tassier-Surine, S.A., 2001, Classic geology of the Burlington Area: Des Moines County, Iowa: Geological Society of Iowa, Guidebook 71, 52 p.
- Witzke, B.J., Bunker, B.J., Anderson, R.R., Artz, J.A., and Tassier-Surine, S.A., 2002, Pleistocene, Mississippian, & Devonian Stratigraphy of the Burlington, Iowa, Area: Iowa Geological Survey Guidebook No. 23, 137 p.
- Witzke, B.J., Anderson, R.R. and Pope, J.P., 2010, Bedrock Geologic Map of Iowa, scale: 1:500,000: Iowa Geological and Water Survey, Open File Digital Map OFM-10-1.