


**SUMMARY REPORT OF THE
SURFICIAL GEOLOGIC MAP OF
THE KEYSTONE SOUTH 7.5' QUADRANGLE,
BENTON COUNTY, IOWA**

**Iowa Geological Survey
Open File Map OFM-20-9
October 2020**

Phil Kerr, Stephanie Tassier-Surine, and Richard Langel

Iowa Geological Survey, IIHR-Hydrosience & Engineering, University of Iowa, Iowa City, Iowa



Iowa Geological Survey, Keith Schilling, State Geologist

Supported in part by the U.S. Geological Survey
Cooperative Agreement Number G19AC00243
National Cooperative Geologic Mapping Program (STATEMAP)
This work was partially supported by a National Science Foundation Award:
Improving Undergraduate STEM Education Grant GP-IMPACT-1600429.



INTRODUCTION

The Keystone South Quadrangle is located in central Iowa on the border of the Iowan Surface landform region, commonly referred to as the Iowan Erosion Surface (IES), and the Southern Iowa Drift Plain (SIDP) (Prior and Korht, 2006). The map area on the IES is dominated by dissected till plains with unnamed reworked sediments covered by thin loess and elongated loess-covered uplands called paha, as well as larger 'erosional inliers' (Ruhe et al., 1968; Prior and Korht, 2006). On the SIDP, the landscape is steeper with thicker packages of loess that can bury paleosols formed in Pre-Illinoian till. The Cedar River to the north and Iowa River to the south have a significant role in shaping the surficial geology in this area. Stratigraphically, this map contains Pre-Illinoian age glacial deposits above Paleozoic carbonates. This area contains no shallow bedrock, with the average depth to bedrock being over 30 m (100 ft). The thickness of Quaternary deposits in the Keystone South Quadrangle can be over 115m (375 ft) in some areas of the quadrangle.

Mapping the Keystone South and Van Horne quadrangles represents the second part of a multi-year program to map the surficial geology of Benton County. This county represents the last corridor along the Cedar River that has not been recently mapped. The counties to the north and east have both been mapped under the STATEMAP program (Quade et al., 1998; Tassier-Surine et al., 2013). This area represents a unique geologic area, being on the southern edge of the IES. Studies conducted during the 1950s and 1960s showed that the often-debated Iowan Glaciation did not occur, e.g. Ruhe et al. (1968). This work indicated that the unique stratigraphy of the area was caused due to an erosional event due to fluvial downcutting. Modern periglacial ideas have challenged the older model of landscape development for the area, and detailed geologic mapping can be used as an important tool for investigating these processes. The majority of the older research's drill cores were collected in the same general vicinity, so a comparison can be made between the two different ideas on landscape development.

PURPOSE

Detailed geologic mapping of the Keystone South quadrangle was completed as part of the Iowa Geological Survey's (IGS) ongoing participation in the United States Geological Survey (USGS) STATEMAP Program. Mapping in the area is part of the IGS Impaired Watershed mapping initiative and provides comprehensive surficial and bedrock geologic information. These maps are the basis for further development of derivative datasets and map products for use by local, county and state decision-makers. In recent years, Iowa's State Mapping Advisory Committee (SMAC) has recommended mapping in areas with environmental concerns related to groundwater quality and land-use planning issues, and/or in rapidly developing areas. The IGS and SMAC recognize the need for maps of varying scales to address the complex environmental issues facing urban and rural Iowans. Mapping in Benton County provides much needed geologic maps in the Middle Cedar River watershed. Many new partners are concentrating efforts on water supply, water quantity and quality, land-use planning, and flood protection studies. Geologic mapping is crucial and foundational for many of these studies, and the project enjoys broad support from the Cedar River Watershed Coalition (CRWC).

From a Quaternary perspective, characterizing the materials deposited during the formation of the IES is a fundamental part of understanding the recent geologic history of both Iowa and the glaciated parts of the Midwest. The Quaternary IES materials are highly variable and poorly consolidated, and therefore

do not provide good groundwater quality protection in shallow rock areas. It is necessary to identify areas of either better consolidated Quaternary materials or bedrock aquitards. Combining the bedrock and surficial map information is allowing stakeholders to address key questions related to shallow rock areas, potential karst issues, aggregate resource potential and protection, and groundwater vulnerability, while achieving the evolving goals of the watershed management plans.

QUATERNARY HISTORY AND REGIONAL SETTING

The map area has a rich and complex Quaternary geologic history punctuated by at least seven periods of glaciation between 2.6 million and 500,000 years ago (Boellstorff, 1978a, b; Hallberg, 1980, 1986). Episodic erosion over the last 500,000 years has led to the destruction of pre-existing glacial landforms associated with Pre-Illinoian glaciations. While this area was not glaciated during the last glacial episode, the Wisconsin, the effects of the paleoclimate can be seen in deposits throughout the area. This period had an intensely cold and windy environment from 21,000 to 16,500 years ago (Walter, 1994; Bettis and Autin, 1997; Mason, 2015). Due to this climate, a periglacial environment prevailed with intensive freeze-thaw action, solifluction, strong winds, thin plant cover, and a host of other periglacial processes (Baker et al., 1986; Walter, 1994; Mason, 2015; Kerr et al., 2019).

During the formation of the IES, soils that had developed on the upland landscape were removed and the underlying Pre-Illinoian till weathering zone was significantly eroded in this area; this resulted in the development of a region-wide colluvial lag deposit referred to as a “stone line.” During this same interval, the Cedar River valley was filled with glacially derived deposits of sand and gravel from the Des Moines Lobe. This outwash provided a local source for wind-derived sediment. The distinctive features of this region are paha and the larger ‘erosional inliers.’ These are isolated landforms and may represent uneroded topographic highs of loess-mantled Pre-Illinoian till. They have a northwest to southeast alignment (McGee, 1891; Ruhe et al., 1968). This orientation seems to coincide with the regional paleoprevailing winds (Muhs et al., 2013; Mason 2015; Kerr et al., 2019). Paha and ‘inliers’ usually have a thick package of wind-blown material, with some locations in the mapping area having over 12 m (40 ft) of loess and fine sand. Where paha are not present, packages of stratified loamy and sandy sediment cover the upland and hill slope positions on the landscape. This material is thought to be the remnants of periglacial processes associated with the formation of the IES. The toeslopes and smaller valleys have thick packages of coarse sand and gravel interspersed with loamy and silty layers. Some valleys, like Prairie Creek, may have over 7 m (24 ft) of this coarse material beneath a mantle of reworked loess. These valley deposits were developed in response to the shifting climate and base levels during the formation of the IES (Bettis and Autin, 1997).

In eastern Iowa, the highly eroded and dissected Pre-Illinoian upland and older terraces are mantled by two Wisconsin loesses. The older Pisgah Formation is thin and includes loess and related slope sediments that have been altered by colluvial hillslope processes. The unit is characterized by the presence of a weakly developed soil recognized as the Farmdale Geosol. It is not uncommon to see the Farmdale developed through the Pisgah Formation and into the underlying older Sangamon Paleosol. The Pisgah loess was most likely deposited on the eastern Iowa landscape from 30,000 to 24,000 years ago (Bettis, 1989) and is typically buried by Peoria Formation loess. The Peoria Formation loess accumulated on stable landsurfaces in eastern Iowa from 25,000 to 21,000 years ago. Peoria Formation eolian materials mantle the upland till units and are present on the Wisconsin outwash terraces. On the uplands, the Peoria Formation is a uniform silt loam; in the valleys the silt commonly grades downward to fine sand. The loess deposits in the mapping area are relatively thin, generally less than 4 meters (12 ft).

Hudson age deposits are associated with fine-grained alluvial, organic, and colluvial sediments and include the DeForest Formation which is subdivided into the Camp Creek, Roberts Creek, and Gunder members. These deposits are present in valleys and upland drainages throughout the map area.

METHODS

Numerous existing sources of geologic information were utilized in the production of the surficial and bedrock geologic maps of the Keystone South Quadrangle including subsurface information, USDA NRCS soil survey data, aerial photography, digital elevation models (DEM's), satellite imagery, landform characteristics, and LiDAR. Where available, engineering borings from public utilities, the Iowa Department of Transportation, and monitoring well records of the USGS were used. Subsurface lithologic and stratigraphic information was mostly derived from analysis of water well cutting samples repositated at the IGS and stored in the IGS online GeoSam database. Over 120 public and private wells in GeoSam, including strip logs, were reviewed for lithology, stratigraphy and locational accuracy, and updated where needed. NRCS digitized soils data (Brown and Highland, 1980) provided information regarding shallow rock areas, helped to guide valley mapping units, and defined slope areas where glacial till is exposed. Quaternary geologists utilized the IGS truck mounted Giddings probe to drill a mix of solid stem and continuous core holes. Twenty new drill holes totaling 363 feet were completed in or near the quadrangle to characterize the Quaternary sediments and establish unit thickness. Samples are being processed for grain-size with all results expected by November, 2020. Laboratory data will be incorporated into the online IGS GeoLab database.

Project geologists combined information from the sources listed above to delineate surficial geologic mapping units at 1:24,000 scale for the Keystone South Quadrangle. IGS mappers used ArcGIS and on-screen digitizing techniques developed during previous STATEMAP projects. The final map entitled 'Surficial Geologic Map of the Keystone South 7.5' Quadrangle, Benton County, Iowa' will be available as a shapefile in the Iowa GEODATA Clearing House (<https://geodata.iowa.gov>), as a PDF file on the IGS Publications website, and will be submitted to the USGS National Geologic Map Database. This Summary Report is also available as a PDF file on the IGS Publications website.

Stratigraphic Framework for East-Central Iowa

The stratigraphic framework for eastern and central Iowa was established nearly 40 years ago (Hallberg, 1980, 1986). Surficial deposits in the map area are composed of six formations: the DeForest, Noah Creek, Peoria, Pisgah, Wolf Creek, and Alburnett formations, as well as unnamed erosion surface sediments. Hudson age deposits associated with fine-grained alluvial, organic, and colluvial sediments include the DeForest Formation which is subdivided into the Camp Creek, Roberts Creek, Gunder, Corrington, and Woden members (Bettis, 1990a; Bettis et al., 1992). The Noah Creek Formation includes coarse sand and gravel associated with outwash from the Des Moines Lobe, as well as coarse to fine-grained fluvial and colluvial deposits associated with local stream and river valleys (Bettis et al., 1996). Unnamed erosion surface sediments consist of reworked till, slopewash, and soliflucted deposits associated with periglacial activity during the late Wisconsin Episode. The Peoria Formation is found across the mapping area and has a silt and sand facies deposited during the Late Wisconsin Episode (McKay, 1979; Bettis et al., 2003). Generally, the sand facies is found downwind (southeast) of the Cedar River, while the silt facies is found on the upland, especially in paha. Thin deposits of Peoria Formation, less than 1 m (3 ft), are found across most of the mapping area. The Pisgah Formation is comprised of loess and colluvium deposited

during the Middle Wisconsin (Bettis, 1990b). Beneath these materials are Pre-Illinoian glacial deposits, which, in Iowa, consist of two formations: the younger Wolf Creek Formation and the Alburnett Formation (Hallberg, 1980). The Wolf Creek Formation is divided into the Winthrop, Aurora, and Hickory Hills members (oldest to youngest). The Alburnett Formation consists of several “undifferentiated” members. This project had 20 new drill holes to assist with the geologic characterization.

Recent studies and mapping indicate that the map area encompasses a complex suite of depositional landforms and sediment sequences related to glaciations, alluviation, subaerial erosion, and wind-blown transport. To map diverse landscapes at 1:24,000 scale, we have selected the most comprehensive mapping strategy- a landform sediment assemblage (LSA) approach. Various landforms are the result of specific processes at work in the geologic system. Landforms typically have similar relief, stratigraphic and sedimentologic characteristics. Recognition of the genetic relationship among landforms and their underlying sediment sequences allows one to generalize and map complex glacial terrains over areas of large extent (Sugden and John, 1976; Eyles and Menzies, 1983). Bettis and others (1999) found that LSA mapping concepts were extremely useful in overcoming the difficulties of mapping in large valleys and noted that LSA's provided a unique opportunity to associate landforms with their underlying sediment packages. Eight landform sediment assemblage units were identified in the map area utilizing aerial imagery, topographic expression, digitized soils, LiDAR, and existing and new subsurface geologic boring information. The following is a description of each landform sediment assemblage listed in order of episode:

HUDSON EPISODE

Qal - Alluvium (DeForest Formation - Undifferentiated) Variable thickness of less than 1 to 5 m (3-16 ft) of very dark gray to brown, noncalcareous or calcareous, stratified silty clay loam, clay loam, loam to sandy loam alluvium and colluvium in stream valleys, on hill slopes, and in closed depressions. May overlie Wolf Creek or Alburnett formation glacial tills, Peoria Formation loess or eolian sand, or Wisconsinan sand and gravel. Associated with low-relief modern floodplains, closed depressions, modern drainageways, or toeslope positions on the landscape. Seasonal high water table and potential for frequent flooding.

Qaf - Alluvial fan (DeForest Formation - Corrington Mbr.) Variable thickness of 2 to 5 m (7-16 ft) of dark brown to yellowish brown, noncalcareous, silt loam to loam with interbedded lenses of fine sand and silts. A pebble lag is commonly found at or near the fan surface. Overlies sand and gravel of the Noah Creek Formation along Prairie and Buckeye creeks. Shallow angled fans occur at the base of low order drainages and colluvial slopes along these drainages.

WISCONSIN EPISODE

Qnw2 - Sand and Gravel (Noah Creek Formation) Generally 2 to 10 m (6-33 ft) of yellowish brown to gray, poorly to well-sorted, massive to well-stratified, coarse to fine feldspathic quartz sand, pebbly sand and gravel with few intervening layers of silty clay. A thin mantle of loess, reworked loess, or fine-grained alluvium may be present. This unit includes silty colluvial deposits derived from the adjacent map units and also encompasses “pre-Gunder Member” deposits that accumulated in low-relief stream valleys during the Wisconsin and Hudson episodes. Seasonal high water table and some potential for flooding.

Qps1 - Loess and Intercalated Eolian Sand (Peoria Formation - silt and/or sand facies) Generally 2 to 5 m (7-16 ft) of yellowish brown to gray, massive, fractured, noncalcareous grading downward to calcareous, silt loam and intercalated fine to medium, well-sorted sand. Sand is most abundant in the lower part of the eolian package. Overlies massive, fractured, loamy glacial till of the Wolf Creek or Alburnett formations with or without the intervening clayey Farmdale/Sangamon Geosol.

Qps1b - Thick Loess and Intercalated Eolian Sand (Peoria Formation - silt and/or sand facies) Generally 5 to 15 m (16-49 ft) of yellowish brown to gray, massive, noncalcareous grading downward to calcareous silt loam and intercalated fine to medium, well-sorted sand. Minimum thickness of 5 m (16 ft) on uplands. Maximum thickness of 2 to 7 m (6-23 ft) of loess occurs on adjacent slopes. Overlies massive, fractured, loamy glacial till of the Wolf Creek or Alburnett formations with or without the intervening clayey Farmdale/Sangamon Geosol.

Qps3 - Loess Shallow to Loamy and Sandy Sediment (Peoria Formation - silt and/or sand facies) Generally 2 – 3 m (7 - 10 ft) of yellowish brown, massive, noncalcareous silt loam and intercalated fine to medium, well sorted, feldspathic quartz sand. Sand, if present, occurs in the lower part of unit. Overlies 0.5 to 1.5 m (2 - 5 ft) of pebbly loam erosion surface sediment which, in turn, overlies eroded, massive, fractured, firm, loamy glacial till of the Wolf Creek or Alburnett formations. Seasonally high water table may occur in this map unit.

Qwa2 - Loamy and Sandy Sediment Shallow to Glacial Till (Unnamed erosion surface sediment) Generally 2 to 8 m (6-26 ft) of yellowish brown to gray, massive to weakly-stratified, well to poorly-sorted loamy, sandy and silty Iowan Erosion Surface sediment. Map unit includes some areas mantled with less than 2 m (7 ft) of Peoria Formation materials (loess and eolian sand). Overlies massive, fractured, firm, glacial till of the Wolf Creek and Alburnett formations. Seasonal high water table may occur in this map unit.

PRE-ILLINOIS EPISODE

Qwa3 - Glacial Till (Wolf Creek or Alburnett formations) - Generally 60 m (200 ft) but can be more than 91 m (300 ft) thick. This mapping unit consists of very dense, massive, fractured, clay loam glacial till of the Wolf Creek or Alburnett formations. Map unit includes some areas mantled with than 2 to 5 m (7-16 ft) of Peoria Formation materials (loess and eolian sand). This mapping unit can be overlain by unnamed erosion surface sediments, colluvium, outwash, loess, eolian sand, or alluvium.

ACKNOWLEDGMENTS

We would like to thank The Miller Family who allowed us access to their property for drilling. Special thanks also to the Benton County Secondary Roads Department for drilling access. A special thanks to the staff of the Benton County Conservation Board for allowing access onto sites and for many discussions of the geology of the area. Drilling was provided by Matthew Streeter of the Iowa Geological Survey (IGS) with the assistance of Sophie Peirce (IGS) and University of Iowa (UI) student Dan Bloch. UI students Nick Johnson, Ellie Biebesheimer, and Dan Bloch assisted with well locations. Special thanks to Kathy Woida of the Natural Resources Conservation Service (retired) and Art Bettis, UI Department of Earth and Environmental Sciences (retired), for assistance with core description and for numerous valuable discussions regarding the geology of the Iowan Erosion Surface. Casey Kohrt and Chris Kahle of the Iowa Department of Natural Resources provided GIS technical help. Administrative support was provided by Suzanne Doershuk, Melissa Eckrich, Teresa Gaffey, Carmen Langel, and Rosemary Tiwari.

REFERENCES

- Baker, R.G., Bettis, E.A.I., D.G., H., 1993. Late Wisconsinan-Early Holocene Riparian Paleoenvironment in Southeastern Iowa. *Geol. Soc. Am. Bull.* 105, 206-212.
- Baker, R.G., Rhodes, R.S., Schwert, D.P., Ashworth, A.C., Frest, T.J., Hallberg, G.R., and Janssens, J.A., 1986. A Full-Glacial Biota from Southeastern Iowa, USA. *Journal of Quaternary Science* 1, 97-101.
- Bettis, E.A.I., 1990. Holocene Alluvial Stratigraphy and Selected Aspects of the Quaternary History of Western Iowa. *Midwest Friends of the Pliocene*.
- Bettis, E.A.I., 1997. Late-Middle and Early-Late Wisconsin Glaciation in North-Central Iowa, *Geological Society of America Abstracts with Programs*.
- Bettis, E.A.I., Hajic, E.R., Quade, D.J., 1999. Geologic Mapping of Large Valleys in Glaciated Regions: The Use of Landform and Landscape Sediment Assemblages for Multi-Use Maps, *Geological Society of America Abstracts with Programs*, 33rd Annual Meeting North-Central Section April.
- Boellstorff, J.D., 1978a. North American Pleistocene Stages Reconsidered in Light of Probable Pliocene-Pleistocene Glaciation. *Science* 202, 305-307.
- Boellstorff, J.D., 1978b. Chronology of Some Late Cenozoic Deposits from the Central United States and the Ice Ages. *Transactions of the Nebraska Academy of Science* VI, 35-49.
- Brown, M.D., Highland, J.D., 1980. Soil Survey of Benton County, Iowa, in: U.S. Department of Agriculture, S.C.S. (Ed.), pp. 180, 155 map sheets.
- Eyles, N., Menzies, J., 1983. *The Subglacial Landsystem, Glacial Geology*. Elsevier, pp. 19-70.
- Hallberg, G.R., 1980. Pleistocene Stratigraphy in East-Central Iowa. *Iowa Geological Survey Technical Information Series No. 10*.
- Hallberg, G.R., 1986. Pre-Wisconsin Glacial Stratigraphy of the Central Plains Region in Iowa, Nebraska, Kansas, and Missouri. *Quaternary Science Reviews- Quaternary Glaciations in the Northern Hemisphere* 5, 11-15.
- Kerr, P.J., Tassier-Surine, S.A., Korht, C.J., 2019. Trends in Eolian Features on the Iowan Erosion Surface, *Geological Society of America Abstracts with Programs*. Geological Society of America, Manhattan, KS.
- Mason, J.A., 2015. Up in the Refrigerator: Geomorphic Response to Periglacial Environments in the Upper Mississippi River Basin, USA. *Geomorphology* 248, 363-381.10.1016/j.geomorph.2015.08.004.
- McGee, W.J., 1891. The Pleistocene History of Northeastern Iowa. U.S. Geological Survey 11th Annual Report, 199-586.
- McKay, E.D., 1979. Wisconsinan Loess Stratigraphy of Illinois, Wisconsinan, Sangamonian, and Illinoian Straigraphy in Central Illinois. *Illinois State Geological Survey*, pp. 95-108.

Muhs, D.R., Bettis, E.A., Roberts, H.M., Harlan, S.S., Paces, J.B., Reynolds, R.L., 2013. Chronology and Provenance of Last-Glacial (Peoria) Loess in Western Iowa and Paleoclimatic Implications. *Quat. Res.* 80, 468-481.10.1016/j.yqres.2013.06.006.

Prior, J.C., Korht, C.J., 2006. *The Landform Regions of Iowa* 2ed. Iowa Geological Survey.

Quade, D.J., Bettis, E.A.I., Ludvigson, G.A., Giglierano, J., Slaughter, M.K., 1998. *Surficial Geologic Materials of Linn County, Iowa*, OFM, 13-4 ed. Iowa Geological Survey Bureau.

Ruhe, R.V., Dietz, W.P., Fenton, T.E., Hall, G.F., 1968. *Iowan Drift Problem, Northeastern Iowa*. State of Iowa- Report of Investigations 7, Iowa City, IA.

Sugden, D.E., John, B.S., 1976. *Glaciers and Landscape*. E. Arnold.

Tassier-Surine, S.A., Quade, D.J., Rowden, R., McKay, R., Liu, H., Giglierano, J., 2013. *Surficial Geology of Black Hawk County, OFM*, 13-4 ed. Iowa Geological Survey.

Walter, J.C., 1994. Ice-Wedge Casts and Relict Polygonal Patterned Ground in North-East Iowa, USA. *Permafrost and Periglacial Processes* 5, 269-282.