



SURFICIAL GEOLOGIC MAP OF THE DES MOINES LOBE OF IOWA Hamilton and Webster Counties	
Geological Survey Bureau Open File Map 2000-1 May 2000	
Prepared by Deborah J. Quade ¹	
James D. Giglierano ¹ E. Arthur Bettis ² Robin L. Wiener ³	
Supported by the U.S. Coological Super	
Cooperative Agreement Number 99HQAG0064 Vational Cooperative Geologic Mapping Program (STATEMAP)	
Energy and Geological Resources Division	
Geological Survey Bureau	
Iowa Department of Natural Resources	
Lyle W. Asell, Interim Director	
ACKNOWLEDGEMENTS	
ns to map's production: Timothy J. Kemmis, Adrian Goetemoeller, Andy Asell, Kathryn Benz, Mary Ellis, ica Overmohle, Shane Smith Iulia F. Tabbut, Stenbania Tassiar Suring, Mark Alexandra, Mark, and C. J.	
chilly J. Bunker, Jackson L. Gilmore, Calvin F. Wolter, Bernard E. Hoyer, Pat J. Lohman, Michael J. Bounk, ec. Drilling was provided under contract by Aquadrill. Assistance in obtaining permission from landowners preces Conservation Service, Fort Dodge Office. A special thanks to the following individuals who graciously	
rilling: Robert Condon, Terry Condon, Edna Schultz, Delmar Estlund, Delbert Dencklau, Ralph Sonnicksen, ern Chalstrom, Margaret Banwell, Russ Slotten, Garret Geisler, Engstrom, Schwendemann, Greg Carlson, Don Anne Kaputska, Ray Allard, Webster County Conservation Board and the Fort Dodge Asphalt Co., Webster Jim Walker, Lyle Anderson. Dennis Doolittle, John VanLangen, Leslie Eggers, Barbara Sylvester, Lawrence	
ilcox, Don Hill, Bruce Berggren, Meryln Hegland, Kevin Larsen, Glen Hodnefield, Lon Crosby and the bard, Hamilton County.	
University of Iowa, 121 Trowbridge Hall, Iowa City, IA 52242-1519 Service, 1617 N. Lake St., Storm Lake, IA 50588	
LEGEND	
Description of Mapping Units	
Hudson Episode	
t (DeForest Formation-Woden Mbr) Generally 2.5 to 11 m of black to very dark gray, calcareous, muck, peat and vium and organic sediments in drained and undrained closed and semi-closed depressions. Overlies a gray, dense loam diamicton (Dows Fm Alden Mbr.) or may overlie Noah Creek Fm. sand and gravel. Low relief depressions and low sags on the landscape. Seasonal high water table.	
Forest Formation-Undifferentiated) Variable thickness of less than 1 m to 5 m of a very dark gray to brown, fied silty clay loam, clay loam, loam to sandy loam alluvium and colluvium in stream valleys, hillslopes and Aay overlie Dows Formation (Morgan or Alden Mbr.), Noah Creek Formation or Pennsylvanian/Mississippian generating modern floadening, and and and an another and an another and the loadeneer	
veoria Formationsand facies) Generally less than 3 m of yellowish brown, massive, calcareous loamy sand to erlie vellowish-brown coarse-grained sand and gravel (Noah Creek Fm.) or it may overlie vellowish to gravish	
eous, stratified loam to silt loam to sandy loam diamicton; textures can be quite variable (Dows FmMorgan ted to a narrow belt on major river valley bottoms or adjacent uplands on the Des Moines Lobe.	
Late Wisconsin Episode ows Formation-Morgan Mbr.) Less than 8 m of yellowish brown, often calcareous and fractured, stratified loam loam diamicton; textures can be quite variable. Overlies gray, calcareous, massive, dense loam diamicton (Dows	
Low to moderate relief, (3 m to 8 m), undulating plains with irregular surface patterns. Seasonal high water table. th lineated ridge forms (Dows Formation-Morgan Mbr.) Less than 3 m of yellowish to grayish brown, often ured, stratified loam to silt loam to sandy loam diamicton; textures can be quite variable. Overlies gray,	
dense loam diamicton (Dows Fm Alden Mbr.). Low relief, (less than 3 m of local relief), slightly undulating surface patterns. Lineated ridge forms (Dows Formation-Morgan Mbr./ Pilot Knob Mbr.) Less than 6 m of en calcareous, stratified loam to silt loam to sandy loam diamicton; textures can be quite variable. Occasionally, primarily of sand and gravel and exhibit evidence of syndepositional collapse (Pilot Knob Mbr.) Overlies gray,	
dense loam diamicton (Dows FmAlden Mbr.). Faint to well-defined lineated ridges oriented transverse to ton till plain. Ridges are moderate to high relief features (3 to 8+ m). Overall landform is a swell and swale al high water table.	
Dows Formation-Morgan Mbr.) Generally 4 to 15 m of yellowish to grayish brown, usually calcareous and oam to silt loam to sandy loam diamicton; textures can be quite variable. Overlies gray, calcareous, massive, n (Dows Fm Alden Mbr.). Moderate to high relief hummocky landform feature exceeds 3-8 m of local relief.	
tannels (Noah Creek Formation) Generally less than 7 m of yellowish brown coarse-grained sand and gravel. eous, massive, dense loam diamicton (Dows Fm Alden Mbr.). In valley positions, it is at the land surface of e modern floodplain it is buried by DeForest Fm. alluvium. Low-relief landforms that may be deposited in broad a loaming learner or in superto abared opicit torreser	
Channels shallow to bedrock (Noah Creek Formation) Generally less than 5 m of yellowish brown coarse- vel. In the northern part of Webster county the unit overlies Mississippian carbonate bedrock. From Fort Dodge es Pennsylvanian bedrock. In valley positions, the unit is at the land surface of older terraces. On the modern	
d by DeForest Fm. alluvium. Low-relief landforms that may be deposited in long, narrow longitudinal terraces or int terraces which overlie bedrock. n (Noah Creek Formation) Thickness can be quite variable from 3 to 12 m of yellowish-brown coarse-grained	
y overlie yellowish to grayish brown usually calcareous, stratified loam to silt loam to sandy loam diamicton; variable (Dows FmMorgan Mbr.) or it may overlie a gray, calcareous, massive, dense loam diamicton (Dows arrow low-relief apron that gently slopes away from the moraine front. Seasonal high water table.	
nt Plain - broad-scale landform features (Dows Formation-Lake Mills Mbr.) Generally less than 3 m of dark ive, calcareous silty clay loam, silt loam overlying a thin basal increment of sand and gravel (<1 m). It may grayish brown usually calcareous, stratified loam to silt loam to sandy loam diamicton; textures can be quite	
-Morgan Mbr.) or it may overlie a gray, calcareous, massive, dense loam diamicton (Dows Fm Alden Mbr.). n with less than 3 m of local relief. Seasonal high water table. Lake Sediments - hummocky isolated ice-walled lake plains-flat-topped hummocks (Dows Formation-Lake	
of sand and gravel (<1 m). It usually overlies yellowish to grayish brown usually calcareous, stratified loam to um diamicton; textures can be quite variable (Dows FmMorgan Mbr. Isolated ice-walled lake plains comprise a orm area. Other hummocks consist of less than 4 m of yellowish to grayish brown usually calcareous, stratified sandy loam diamicton (Dows FmMorgan Mbr.). Overlies a gray calcareous measing dance to the stratified	
Mbr.). Moderate to high relief hummocky landform area with 3 to 8+ m of local relief. ment Plain - broad-scale landform features with lineated ridge forms (Dows Formation- Lake Mills Generally less than 3 m of dark grayish brown, massive, calcareous silty clay loam silt loam overlying a thin	
ind and gravel (<1 m). It may overlie yellowish to grayish brown usually calcareous, stratified loam to silt loam cton; textures can be quite variable (Dows FmMorgan Mbr.) or it may overlie a gray, calcareous, massive, on (Dows Fm Alden Mbr.). Low relief broad plain with less than 3 m of local relief. Faint lineated ridges placier flow are inset on till plain adjacent to major stream and river valleys. Ridges are lower relief features	
orm is an undulating plain. Low relief (<3 m) on lineated ridges. Seasonal high water table.	
Complex (Dows Formation-Pilot Knob Mbr./ Morgan Mbr.) Greater than 3 m and less than 15 m of yellowish ous and fractured, stratified sand and gravel with interbedded stratified loam diamicton. Collapse features are depressions and sags on upland surfaces, the sand and gravel may be buried by DeForest Fm -Woden Mbr	
eous, massive, dense loam diamicton (Dows Fm Alden Mbr.). Moderate to high relief (3 to 8+ m) hummocks, s and associated linked drainages on upland surfaces.	
al Complex with lineated ridge forms (Dows Formation-Pilot Knob Mbr/ Morgan Mbr.) Greater than 4 m and vellowish brown, often calcareous and fractured, stratified sand and gravel with interbedded stratified loam sh to grayish brown, often calcareous and fractured, stratified loam to silt loam to sandy loam diamicton; textures . In depressions and sags on upland surfaces, the sand and gravel may be buried by DeForest FmWoden Mbr.	
eous, massive, dense loam diamicton (Dows Fm Alden Mbr.). Faint to well-defined lineated ridges oriented flow are inset on till plain. Ridges are moderate to high relief features (3 to 8+ m). Occasionally, these ridges sand and gravel and exhibit evidence of syndepositional collapse (Pilot Knob Mbr.). Overall landform can be and swale topography. Moderate to high relief (3 to 8+ m) on lineated ridges and hummocks. Seasonal high water	
nian Bedrock(Cherokee Group)/Mississippian Bedrock (St. Louis and Pella Formations) Pennsylvanian	
of deltaic sediments associated with the Cherokee Group. Lithologies are extremely variable. Major lithologies sandstone with minor coal seams. <i>Mississippian</i> Composed of marine limestones, sandstones and shales. ng the Des Moines River valley north of Fort Dodge. In the Fort Dodge vicinity some outcrops of Jurassic age stones are present.	
rries Gypsum and limestone quarries; and sand, gravel, and clay pits. Extent mapped as shown in county soil	
najor land filling. Fill associated with railroad grades, highway grades and land leveling. Variable in texture to sandy to concrete rubble. Extent manned as shown in county soil surrays	
-, a racere, zatent mapped as snown in county son surveys.	